



## A new species of leopard frog (Anura: Ranidae) from the urban northeastern US

Catherine E. Newman<sup>a,\*</sup>, Jeremy A. Feinberg<sup>b</sup>, Leslie J. Rissler<sup>c</sup>, Joanna Burger<sup>b</sup>, H. Bradley Shaffer<sup>a,d,1</sup>

<sup>a</sup> Department of Evolution and Ecology, University of California, Davis, CA 95616, USA

<sup>b</sup> Graduate Program in Ecology & Evolution, Department of Ecology, Evolution, and Natural Resources, Rutgers University, New Brunswick, NJ 08901, USA

<sup>c</sup> Department of Biological Sciences, University of Alabama, Tuscaloosa, AL 35487, USA

<sup>d</sup> Center for Population Biology, University of California, Davis, CA 95616, USA

### ARTICLE INFO

#### Article history:

Received 25 August 2011

Revised 20 January 2012

Accepted 22 January 2012

Available online 2 February 2012

#### Keywords:

*Rana pipiens*

*Rana sphenocephala*

*Lithobates*

Urban ecology

Amphibian decline

Species delimitation

### ABSTRACT

Past confusion about leopard frog (genus *Rana*) species composition in the Tri-State area of the US that includes New York (NY), New Jersey (NJ), and Connecticut (CT) has hindered conservation and management efforts, especially where populations are declining or imperiled. We use nuclear and mitochondrial genetic data to clarify the identification and distribution of leopard frog species in this region. We focus on four problematic frog populations of uncertain species affiliation in northern NJ, southeastern mainland NY, and Staten Island to test the following hypotheses: (1) they are conspecific with *Rana sphenocephala* or *R. pipiens*, (2) they are hybrids between *R. sphenocephala* and *R. pipiens*, or (3) they represent one or more previously undescribed cryptic taxa. Bayesian phylogenetic and cluster analyses revealed that the four unknown populations collectively form a novel genetic lineage, which represents a previously undescribed cryptic leopard frog species, *Rana* sp. nov. Statistical support for *R. sp. nov.* was strong in both the Bayesian (pp = 1.0) and maximum-likelihood (bootstrap = 99) phylogenetic analyses as well as the Structure cluster analyses. While our data support recognition of *R. sp. nov.* as a novel species, we recommend further study including fine-scaled sampling and ecological, behavioral, call, and morphological analyses before it is formally described.

© 2012 Elsevier Inc. All rights reserved.

### 1. Introduction

Leopard frogs of the *Rana pipiens* (= *Lithobates pipiens*) complex are widespread and common throughout much of the United States, but species delimitation and the associated taxonomy of the group have been challenging and contentious (Brown, 1973; Pace, 1974; Moore, 1975; Brown et al., 1977, 1990; Zug et al., 1982; Hillis, 1988; Frost et al., 2006, 2008, 2009; Pauly et al., 2009). While studies of range-wide phylogeography and systematics at the genus and species level are common (e.g., Pace, 1974; Hillis et al., 1983; Pytel, 1986; Hoffman and Blouin, 2004; Hillis and Wilcox, 2005; Oláh-Hemmings et al., 2010; Newman and Rissler, 2011), relatively little attention has been focused on taxonomic status and conservation needs of local or regional populations or

subspecies (but see Di Candia and Routman, 2007; Hekkala et al., 2011). As is true for any group, appropriate conservation measures cannot be identified and implemented in the face of uncertain taxonomy (Köhler et al., 2005).


The species composition of leopard frogs in parts of the mid-Atlantic and northeastern US—hereafter the Tri-State area, including New Jersey (NJ), New York (NY), and Connecticut (CT)—has been questioned by biologists over the past several decades (Kaufeld, 1937; Yeaton, 1968; Schlauch, 1971; Pace, 1974; Klemens et al., 1987; Klemens, 1993). Currently, two species are recognized in the region (Conant and Collins, 1998). *Rana pipiens*, the northern leopard frog, is widely distributed across New England and the Great Lakes region, including the western two-thirds of CT and central and northern NY. From NJ, Long Island (NY), and southern mainland NY to the south, it is replaced by *R. sphenocephala* (= *L. sphenocephalus*), the southern leopard frog. While natural history collection data suggest the two species have a narrow zone of overlap in southern NY (Fig. 1), no area of sympatry has been directly identified. Some earlier studies based on morphological data suggested the possibility of intergradation (Schlauch, 1971), whereas others speculatively discussed a putative third species in this region (Kauffeld, 1937; Klemens, 1993).

Although widespread and often common at the continental scale (Fig. 1), leopard frog populations have been severely declining in certain regions, resulting in extirpation from some portions

\* Corresponding author. Present address: Department of Biological Sciences, Museum of Natural Science, 119 Foster Hall, Louisiana State University, Baton Rouge, LA 70803, USA. Fax: +1 225 578 3075.

E-mail addresses: [cnewm15@lsu.edu](mailto:cnewm15@lsu.edu) (C.E. Newman), [jerfein@eden.rutgers.edu](mailto:jerfein@eden.rutgers.edu) (J.A. Feinberg), [rissler@as.ua.edu](mailto:rissler@as.ua.edu) (L.J. Rissler), [burger@biology.rutgers.edu](mailto:burger@biology.rutgers.edu) (J. Burger), [brad.shaffer@ucla.edu](mailto:brad.shaffer@ucla.edu) (H.B. Shaffer).

<sup>1</sup> Present addresses: Department of Ecology and Evolutionary Biology, University of California, 621 Charles E. Young Dr. South, Los Angeles, CA 90095-1606, USA, & La Kretz Center for California Conservation Science, Institute of the Environment and Sustainability, University of California, La Kretz Hall, Suite 300, 619 Charles E. Young Dr. South, Los Angeles, CA 90095-14966, USA.



**Fig. 1.** Range maps for *Rana pipiens* (light gray shading) and *R. sphenoccephala* (dark gray shading) in the US. Black indicates range overlap. Inset: sampling localities for genetic analyses. Numbers correspond to Table 1. Green: *R. sphenoccephala* range, blue: *R. pipiens* range, dark gray: range overlap. Red oval contains the four focal populations in this study. NY: New York, PA: Pennsylvania, NJ: New Jersey, CT: Connecticut, MA: Massachusetts, SI: Staten Island, LI: Long Island. Range maps were downloaded as ESRI shapefiles from the IUCN Red List spatial data collection (2011). Colors are available in the online version. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

of their historical range (Lannoo, 2005), including coastal regions and islands north and east of Long Island, NY (Ditmars, n.d.; Latham, 1971; Klemens, 1993; Feinberg, et al., unpublished data). Leopard frogs are also believed to be extirpated from highly developed areas including Long Island, NY (Kiviat, 2010; Feinberg et al., unpublished data); New Haven, CT; and Providence, Rhode Island (Klemens, 1993). While the exact causes of these declines are unclear, environmental pesticides and endocrine disruptors (Hayes et al., 2003; Lannoo, 2008), disease (Carey et al., 1999; Greer et al., 2005; Davis et al., 2007; Searle et al., 2011), habitat loss and alteration (Lannoo, 2005), and over-harvesting for use as laboratory specimens (Hillis, 1988; Klemens, 1993; Lannoo, 2005) have all been identified as contributing factors, particularly regarding *R. pipiens*. *Rana sphenoccephala*, in contrast, remains relatively abundant throughout most of its range to the south, including coastal islands south of Long Island. However, near its northern range limit, it is listed as a Species of Special Concern in NY (NY Department of Environmental Conservation) and as endangered in Pennsylvania (PA) (Pennsylvania Fish and Boat Commission).

To gain a better understanding of the status and distributions of leopard frog populations in the Tri-State area, we analyzed mitochondrial and nuclear gene sequences from four focal populations of unknown leopard frog species composition in northern NJ, southeastern mainland NY (two populations), and Staten Island, NY (one of the five boroughs of New York City). Direct observations by one of us (JAF) showed that these four populations exhibited several unique characteristics, including an advertisement call distinct from both *R. pipiens* and *R. sphenoccephala*. We also analyzed

three CT populations from localities within the traditionally accepted geographic range of *R. pipiens*. We evaluated three possible interpretations of the status of leopard frogs in the Tri-State area: (1) the four focal populations are conspecific with either *R. pipiens* or *R. sphenoccephala*, (2) the populations are hybrids between *R. pipiens* and *R. sphenoccephala*, or (3) the populations represent a previously undescribed leopard frog lineage distinct from *R. pipiens* and *R. sphenoccephala*.

## 2. Materials and methods

### 2.1. Study area and sample collection

Our study region was focused on the Tri-State area of the northeastern US, including NY, NJ, and CT—a total area of roughly 40,000 km<sup>2</sup> (Fig. 1). The region includes an area of putative range overlap between *R. sphenoccephala* and *R. pipiens* according to range maps downloaded from the IUCN [IUCN Red List of Threatened Species 2011.1 (<http://www.iucnredlist.org>)]. Our study included four focal populations of unknown leopard frog species composition: Great Swamp (NJ), Staten Island (NY), Putnam County (NY), and Orange County (NY) (Fig. 1). The Great Swamp and Staten Island sites fall within the geographic range of *R. sphenoccephala* and outside the range of *R. pipiens*, whereas the Putnam and Orange sites fall in the overlap zone of the two species' ranges. Leopard frog species composition in CT has also been questioned (Klemens, 1993), so we collected samples from three sites across CT to include in the analyses (Fig. 1).

Toe clips were taken from 3 to 10 individual frogs at each of the four focal sites and three populations in CT, as well as control sites for *R. sphenoccephala* in southern NJ and *R. pipiens* in northeastern mainland NY (Fig. 1; Table 1). In addition, three morphologically ambiguous specimens from Long Island were included to determine if they represented an isolated relict population of leopard frogs, or if they were instead the pickerel frog *R. palustris* (= *L. palustris*) (Table 1). Three CT *R. palustris* specimens from the Yale Peabody Museum were also included as reference samples. Tissues were stored in 98% ethanol, and source frogs were measured, photographed, and subsequently released, or collected as vouchers to be deposited in either the Yale Peabody Museum or the University of Alabama Herpetological Collection.

## 2.2. DNA extraction and gene amplification

Genomic DNA was extracted at the University of California, Davis, using a standard salt extraction protocol. We amplified the ND2 and 12S–16S regions of the mitochondrial genome, including the intervening tRNA-Valine and partial flanking tRNA-Tryptophan sequences, for a total of 1444 bp. We also amplified the neurotrophin-3 (NTF3, 599 bp), tyrosinase (Tyr, 557–585 bp), Rag-1 (647–683 bp), seven-in-absentia (SIA, 362–393 bp), and chemokine receptor 4 (CXCR4, 550 bp) regions of the nuclear genome. All primer references and sequences are provided in Supplementary Table S1.

PCR amplification was performed in 18  $\mu$ L reactions, consisting of 1.5  $\mu$ L PCR Buffer II (10X, Applied Biosystems), 2.4  $\mu$ L MgCl<sub>2</sub> (25 mM), 0.6  $\mu$ L each primer (5 mM), 2.4  $\mu$ L dNTP solution (5 mM), 1U AmpliTaq (Applied Biosystems), and 10–30 ng genomic DNA. All gene regions except 12S–16S were amplified using the following PCR protocol: initial denaturation at 95° for 1 min.; 38 cycles of 94° for 30 s, 63–65° (see Supplementary Table S1) for 45 s, 72° for 1 min.; and a final extension at 72° for 10 min. The amplification protocol for the 12S–16S region was as follows: initial denaturation at 94° for 30 s; 35 cycles of 94° for 45 s, 52° for 30 s, 72° for 1 min. and a final extension at 72° for 7 min. PCR products were sequenced in the forward and reverse directions at Beckman Coulter Genomics (Danvers, MA, USA). Contigs were assembled in Geneious v.5.3.6 (Drummond et al., 2011). Sequence fragments were trimmed to minimize missing data.

## 2.3. Mitochondrial sequence analysis

The 12S–16S and ND2 sequence fragments with associated tRNA fragments were concatenated and aligned using ClustalW

in Geneious and manually adjusted. All sequences were uploaded to GenBank (see Supplementary Table S2 for accession numbers). Bayesian analyses were conducted in MrBayes 3.1 (Huelsenbeck and Ronquist, 2001; Ronquist and Huelsenbeck, 2003) with five partitions: 12S plus tRNA-Val, 16S, and each of the three ND2 codon positions. Based on output from jModelTest v.0.1.1 (Guindon and Gascuel, 2003; Posada, 2008) and convergence analyses of trial runs (data not shown), the 12S and ND2 partitions were assigned a GTR model of evolution, and the 16S partition was assigned an HKY model of evolution. The 12S partition allowed across-site rate variation under a gamma distribution, and rates were allowed to vary among partitions. Bayesian analyses were run with random starting trees, two simultaneous runs of 10 million generations, and sampling from the posterior distribution of trees every 5000 generations. Tracer v.1.4.1 (Rambaut and Drummond, 2007) was used to assess convergence and to determine appropriate burn-in. The first 25% of samples were omitted as burn-in. Nodal support was further assessed with a maximum-likelihood (ML) analysis in RaxML v.7.0.3 (Stamatakis, 2006; Stamatakis et al., 2008) with 1000 bootstraps. *Rana clamitans* sequences (DQ347036, 12S–16S; AY206480, ND2) were downloaded from GenBank and used as an outgroup. Tajima's *D* and Fu's *F<sub>s</sub>* were calculated in Arlequin v.3.5 (Excoffier and Lischer, 2010) to test for selection.

## 2.4. Nuclear sequence analysis

For each locus, sequences were aligned using ClustalW in Geneious and manually adjusted, and sequences were uploaded to GenBank (Supplementary Table S2). Phylogenies were reconstructed for each locus individually and for the concatenated data set using unphased sequences (see below) in MrBayes. For the individual gene trees, models of evolution, based on jModelTest output and preliminary runs (data not shown), were as follows: HKY for CXCR4; HKY + G for NTF3, Rag-1, and Tyr; and JC for SIA. The concatenated data set was partitioned by locus, but a consistent lack of convergence suggested that this model was inappropriate for our data (results not shown). Bayesian analyses were thus run on the entire, unpartitioned nuclear data set, with an HKY model of evolution [based on jModelTest and trial runs (data not shown)]. All analyses were run for 10 million generations and sampled every 5000 generations. Convergence was assessed in Tracer, and the first 25% of samples were omitted as burn-in. Nodal support was further assessed with 1000 ML bootstraps in RaxML. Tests for selection were done in Arlequin, using Tajima's *D* and Fu's *F<sub>s</sub>* statistics.

**Table 1**


Specimens used in genetic analyses. GSNWR = great swamp national wildlife refuge, BRSP = bass river state park. More specific locality information is available from authors. Sample IDs are listed in Supplementary Table S1.

Population	Sample size	Species ( <i>a priori</i> )	Map code	County	State
<i>Focal unknown populations</i>					
Staten Island	6	Unknown	1	Richmond	New York
GSNWR	5	Unknown	2	Morris	New Jersey
Putnam	5	Unknown	3	Putnam	New York
Orange	3	Unknown	4	Orange	New York
<i>Unknown populations in connecticut</i>					
Hartford	5	Unknown	5	Hartford	Connecticut
Middlesex	5	Unknown	6	Middlesex	Connecticut
Litchfield	10	Unknown	7	Litchfield	Connecticut
<i>Unknown populations on long Island</i>					
Eastport	3	Unknown	10	Suffolk	New York
<i>Control populations (known species)</i>					
BRSP	5	<i>R. sphenoccephala</i>	8	Burlington	New Jersey
Saratoga	6	<i>R. pipiens</i>	9	Saratoga	New York
Litchfield	1	<i>R. palustris</i>	7	Litchfield	Connecticut
New London	1	<i>R. palustris</i>	11	New London	Connecticut
Fairfield	1	<i>R. palustris</i>	12	Fairfield	Connecticut

To test our hypotheses concerning the status of the four unknown populations, we used a Bayesian approach implemented in Structure v.2.3.3 (Pritchard et al., 2000; Falush et al., 2003) with an allelic data set (6% missing data) generated from our nuclear sequence data. We used the software Phase v.2.1 (Stephens et al., 2001; Stephens and Donnelly, 2003) to infer haplotypes for each locus in the five-locus sequence data set using a Bayesian algorithm. Each allele represented a single haplotype. Input files for

Phase were generated from alignment nexus files using a Perl script (RC Thomson, unpublished).

Structure was used to determine the number of genetically distinct clusters ( $K$ ) of samples. We implemented the admixture model (Pritchard et al., 2000), assumed correlation of allele frequencies among clusters (Falush et al., 2003), and assumed no other *a priori* population information. We tested values of  $K$  from 1 to 10. For each  $K$ , 20 iterations were run, each consisting of 100,000



**Fig. 2.** Bayesian phylogeny for concatenated mtDNA (12S–16S and ND2). Nodal support: Bayesian posterior probabilities/maximum-likelihood bootstrap values. Tip labels correspond to Supplementary Table S2. Clade symbols correspond to Fig. 1.

generations after a burn-in of 10,000 generations. The appropriate value of  $K$  was determined by assessing the posterior probabilities (Pritchard et al., 2000) and  $\Delta K$  values following Evanno et al. (2005). An individual was considered of mixed ancestry if its cluster membership probability  $q$  was between 0.10 and 0.90 (Vähä and Primmer, 2005).

For both mitochondrial and nuclear loci, measures of sequence divergence (uncorrected  $p$ ), nucleotide diversity ( $\pi$ ) and haplotype diversity (Hd) were determined at the species level using either DnaSP v.5.10.01 (Librado and Rozas, 2009) or Arlequin. Pairwise  $F_{ST}$  values were calculated in Arlequin from the concatenated, phased nuclear sequence data set.

### 3. Results

#### 3.1. mtDNA phylogenetic analyses

The concatenated mtDNA data set consisted of 1461 bp and 15 unique haplotypes. Bayesian analyses of mtDNA revealed four distinct clades, three of which correspond to the known species *R. sphenoccephala*, *R. pipiens*, and *R. palustris* (Fig. 2). The three samples from Long Island fell out with the *R. palustris* reference samples, rejecting the hypothesis that those frogs represented a relict population of leopard frogs on Long Island. All specimens from the four focal populations and three of five specimens from Middlesex, CT, formed a clade (hereafter *Rana* sp. nov.) distinct from *R. sphenoccephala*, *R. pipiens*, and *R. palustris*. All other CT specimens grouped with *R. pipiens*. All four clades were strongly supported with Bayesian posterior probabilities (all 1.0) and ML bootstraps (all  $\geq 99$ ). Importantly, the sister group to the *R. sp.*

nov. clade, with reasonably strong statistical support (Bayesian posterior probability = 1.0, ML bootstrap support = 0.78) is the pickerel frog *R. palustris* rather than *R. pipiens* or *R. sphenoccephala*. Species-level  $\pi$  and Hd values are listed in Table 2. Pairwise sequence divergence between *R. sp. nov.* and the three recognized species were 6.79% (*R. palustris*), 11.0% (*R. sphenoccephala*), and 12.5% (*R. pipiens*), and pairwise divergence between the latter three described species ranged from 11.1% to 13.4% (Table 3). These data indicate that a differentiated lineage, distinct from *R. sphenoccephala*, *R. pipiens*, and *R. palustris*, occurs in the region and may represent a previously unrecognized species if additional data confirm these mtDNA results.

The three outlier specimens from Middlesex, CT, are more difficult to interpret. Phylogenetically, they fell in the *R. sp. nov.* clade, but their geographic location substantially farther east than all other *R. sp. nov.* samples and, more importantly, their sympatry with *R. pipiens* at the same site made us question whether they represent a natural population of *R. sp. nov.* farther east than expected or human-mediated introductions. Given this uncertainty, we omitted these individuals from population genetic calculations, pending future sampling in CT, particularly the region between Middlesex County and the closest known *R. sp. nov.* population roughly 95 km due west in Putnam County, NY.

#### 3.2. Nuclear phylogenetic analyses

Aligned sequence lengths for nuclear loci were 550 bp (CXCR4), 599 bp (NTF3), 683 bp (Rag-1), 393 bp (SIA), and 585 bp (Tyr). The concatenated data set consisted of 2810 bp of aligned, trimmed sequence. The number of variable sites for each locus ranged from 10 to 30 (Table 2). Species-level  $\pi$  and Hd values are listed in Table 2,

**Table 2**

Species-level general descriptive statistics. Length = aligned sequence length (bp), #VS = number of variable sites, Hd = haplotype diversity,  $\pi$  = nucleotide diversity,  $\theta_\pi$  = number of pairwise nucleotide differences,  $\theta_S$  = Watterson's estimator of genetic diversity.

	Descriptive statistics					Tests of neutrality				
	Length	#VS	Hd	$\pi$	$\theta_\pi$	$\theta_S$	Tajima's $D$	$P$ -value	Fu's $F_S$	$P$ -value
mtDNA	1444	309								
<i>R. sphenoccephala</i>			0.833	0.0013	1.833	1.636	1.09	0.854	0.006	0.292
<i>R. palustris</i>			0.6	0.0048	1.2	0.876	1.753	0.988	1.938	0.798
<i>R. pipiens</i>			0.786	0.0008	7.029	5.003	1.505	0.943	3.215	0.9
<i>Rana</i> sp. nov.			0.582	0.0004	0.619	2.018	0.262	0.688	0.235	0.62
CXCR4	550	13								
<i>R. sphenoccephala</i>			0.429	0.0016	1.333	1.414	-0.219	0.401	1.056	0.748
<i>R. palustris</i>			0.378	0.0021	0.956	1.414	-1.245	0.112	0.39	0.494
<i>R. pipiens</i>			0.633	0.0017	1.228	1.138	0.196	0.62	0.381	0.613
<i>Rana</i> sp. nov.			0.508	0.0012	0.617	0.94	-0.822	0.237	0.688	0.605
NTF3	599	10								
<i>R. sphenoccephala</i>			0.533	0.0009	0.533	0.353	1.302	0.95	1.029	0.635
<i>R. palustris</i>			0.485	0.0012	0.485	0.331	1.066	0.886	1.003	0.569
<i>R. pipiens</i>			0.518	0.0008	1	1.593	-1.001	0.169	-1.128	0.284
<i>Rana</i> sp. nov.			0.514	0.0009	0.511	0.232	1.688	0.978	1.886	0.771
Rag-1	683	20								
<i>R. sphenoccephala</i>			0.778	0.0025	1.711	1.414	0.807	0.785	0.251	0.538
<i>R. palustris</i>			0.8	0.001	1.848	1.656	0.429	0.696	0.737	0.677
<i>R. pipiens</i>			0.273	0.0043	0.594	0.683	-0.273	0.459	2.289	0.847
<i>Rana</i> sp. nov.			0.605	0.0018	1.533	1.394	0.263	0.653	0.94	0.716
SIA	393	19								
<i>R. sphenoccephala</i>			0.429	0.0065	2.8	2.121	1.325	0.907	5.13	0.983
<i>R. palustris</i>			0.439	0.0004	0.47	0.662	-0.85	0.246	-0.725	0.097
<i>R. pipiens</i>			0.165	0.0012	0.168	0.455	-1.148	0.089	-1.722	0.025
<i>Rana</i> sp. nov.			0.163	0.0019	1.232	1.609	-0.637	0.284	1.349	0.793
Tyrosinase	585	30								
<i>R. sphenoccephala</i>			0.929	0.0085	4.689	4.595	0.093	0.551	-0.892	0.262
<i>R. palustris</i>			0	0.0053	0	0	0	1	0	-
<i>R. pipiens</i>			0.803	0	3.065	2.048	1.412	0.926	2.73	0.886
<i>Rana</i> sp. nov.			0.767	0.0075	4.093	2.759	1.464	0.94	4.154	0.949



**Table 3**Intraspecific and pairwise percent sequence divergence (uncorrected *p*), %SD, for mtDNA.

	Intraspecific (%)	Pairwise		
		<i>R. sphenoccephala</i> (%)	<i>R. palustris</i> (%)	<i>R. pipiens</i> (%)
<i>R. sphenoccephala</i>	0.11	–		
<i>R. palustris</i>	0.08	11.1	–	
<i>R. pipiens</i>	0.43	13.4	12.8	–
<i>Rana</i> sp. nov.	0.04	11.0	6.8	12.5

**Table 4**Species-level pairwise  $F_{ST}$ , based on phased nuDNA.

	<i>R. sphenoccephala</i>	<i>R. palustris</i>	<i>R. pipiens</i>
<i>R. sphenoccephala</i>	–		
<i>R. palustris</i>	0.661	–	
<i>R. pipiens</i>	0.463	0.627	–
<i>Rana</i> sp. nov.	0.423	0.695	0.536

and pairwise  $F_{ST}$  in Table 4. Tajima's *D* and Fu's  $F_S$  tests for selection were non-significant for all loci (Table 2), indicating that all sampled loci were selectively neutral.

Analyses of individual nuclear loci (Fig. 3) revealed varying degrees of support for the four species recovered in the mtDNA analysis (Fig. 2). Monophyly of *R. palustris* was strongly supported by four loci, *R. pipiens* by two loci, and *Rana* sp. nov. by one locus. None of the loci supported a monophyletic *R. sphenoccephala*. Importantly, none of the loci recovered strong clade support for non-monophyly of any of the species. In other words, no strongly supported clade contained individuals of multiple species.

Bayesian analysis of the concatenated data set recovered three strongly supported clades corresponding to the three known species (*R. sphenoccephala*, *R. pipiens*, *R. palustris*) (Fig. 4), although their interrelationships were unresolved. The remainder of the samples—those that formed the *R. sp. nov.* mtDNA clade—constituted an unresolved collection of samples that were excluded from all three currently recognized species. While we acknowledge the problems associated with phylogenetic analyses of concatenated nuclear data sets (e.g., Kubatko and Degnan, 2007), we emphasize the concordance among the delimitations in our mitochondrial (Fig. 2) and concatenated nuclear (Fig. 4) phylogenies, as well as the Structure analysis (Fig. 5, see below).

The number of inferred haplotypes per locus ranged from 10 to 19. Bayesian cluster analyses in Structure recovered four clusters ( $\ln L = -504.0$ ,  $\Delta K = 224.13$ ) consistent with the phylogenetic analyses (Fig. 5). As in the mtDNA analyses, *R. sphenoccephala* grouped together in one cluster, *R. palustris* reference samples grouped with the Long Island specimens in a second cluster, all specimens from CT except three from Middlesex grouped with *R. pipiens* controls, and all specimens from the four focal populations grouped with three from Middlesex, CT, in a fourth cluster (*R. sp. nov.*). The three specimens from Middlesex, CT, that clustered with *R. sp. nov.* are the same three that clustered with this group in the mtDNA sequence analyses. Cluster membership values of samples, *q*, ranged from 0.922 to 0.992. None of the samples were of admixed ancestry.

## 4. Discussion

### 4.1. Taxonomic status and geographic distribution of *Rana* sp. nov.

Our data strongly support the recognition of three evolutionary lineages of leopard frogs in the Tri-State area, with the four focal populations collectively forming a new, previously undescribed leopard frog species (*R. sp. nov.*). Phylogenetic and cluster analyses


revealed the four unknown populations to be a distinct group from all locally occurring, recognized leopard frog species, rejecting the hypotheses that those populations are conspecific with one or more of the known species or that they are admixed, intergrade populations. Mitochondrial pairwise sequence divergences between *R. sp. nov.* and the currently recognized species ranged from 6.79% to 12.9%, consistent with or greater than divergence estimates among other ranid species (Jaeger et al., 2001; Shaffer et al., 2004; Di Candia and Routman, 2007; Funk et al., 2008; Oláh-Hemmings et al., 2010). These high levels of divergence strongly suggest a lack of gene flow between *R. sp. nov.* populations and other leopard frog species, and cluster analysis indicated that none of the samples were of admixed ancestry.

Empirical methods for species delimitation (Sites and Marshall, 2004) could potentially add support to our conclusions. In addition, new methods have recently become available that use Bayesian analyses of multilocus sequence data to concurrently estimate the species tree and delimit species (O'Meara, 2010; Niemiller et al., 2011). We argue that such analyses are not necessary in our case, however, because species delimitation is relatively straightforward given the data herein. The older species are, the more time they have had to accumulate various evidences of lineage divergence, such as diagnosable morphological characters, reproductive isolation, or reciprocal monophyly (de Queiroz, 2007; Shaffer and Thomson, 2007). In our study, genetic data suggest monophyly of each of the four species, and the sympatry of *R. pipiens* and *Rana* sp. nov. in Middlesex, CT, suggests some extent of reproductive isolation between the two. Together, reciprocal monophyly and reproductive isolation strongly indicate the reality of independently evolving lineages, which we designate as distinct species.


Based on our current, relatively sparse sampling, *R. sp. nov.* is restricted to northern NJ, extreme southeastern mainland NY, and Staten Island (Fig. 1), although range limits may extend as far as CT and northeastern PA (Pace, 1974). Three samples from Middlesex County, CT, suggest that the range potentially extends into the western half of CT, where *R. sp. nov.* is currently sympatric with *R. pipiens*. Additional sampling in western CT should help to clarify the range extent of *R. sp. nov.* However, we reiterate that our results show no evidence of hybridization between *R. sp. nov.* and either of the other two leopard frog species in the region, including central CT where *R. sp. nov.* and *R. pipiens* occur in sympatry, suggesting some level of reproductive isolation.

### 4.2. Conservation implications and recommendations

The geographic extent of *R. sp. nov.* is limited to a small portion of NJ, NY, and possibly CT and PA (Fig. 1). This northeastern endemic distribution is concordant with few other amphibian taxa (but see *Pseudacris kalmi*; Lemmon et al., 2007) and presents a unique situation compared to more “standard” amphibian phylogeographic patterns (Rissler and Smith, 2010). Pending additional field sampling, the recognition of a distinct, geographically-restricted species suggests that conservation needs may be high, particularly in light of the tremendous human population density in this region



**Fig. 3.** Bayesian phylogenies for individual nuDNA loci: (a) CXCR4, (b) NTF3, (c) Rag-1, (d) SIA, (e) Tyr. Nodal support: Bayesian posterior probabilities/maximum-likelihood bootstrap values. Outgroup root (*R. catesbeiana*) was removed for diagram simplicity. Tip labels correspond to [Supplementary Table S2](#). Clade symbols correspond to [Fig. 1](#). Colors correspond to inferred species: *R. sphenocephala* (green), *R. pipiens* (blue), *R. palustris* (orange), *Rana* sp. nov. (red). Colors are available in the online version. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)



**Fig. 4.** Bayesian phylogeny for concatenated nuDNA (CXCR4, NTF3, Rag-1, SIA, Tyr). Nodal support: Bayesian posterior probabilities. Tip labels correspond to Supplementary Table S2.






Fig. 5. Structure bar plot based on nuDNA. Population numbers are in parentheses under the text label and correspond to Table 1 and Fig. 1. Focal populations are marked with asterisks.

and epidemic declines and extirpations from mainland and coastal regions of the Tri-State area, including Long Island (Feinberg et al., unpublished data), an area once considered a regional stronghold for leopard frogs (Schlauch, 1978).

*Rana sphenoccephala* is currently (as of 2011) listed as a Species of Special Concern in NY; it is not listed in NJ. *Rana pipiens* is not a listed species in NY and is not known to be present in NJ. Our genetic data demonstrate that all of the leopard frogs collected in southern mainland NY for this study were *R. sp. nov.*, rather than *R. sphenoccephala*. Staten Island and the two populations in southern mainland NY (Orange, Putnam) are the only known extant putative *R. sphenoccephala* populations in NY, suggesting that southern leopard frogs do not occur in NY, although information gaps remain regarding Long Island. Furthermore, *R. sphenoccephala* is currently believed to be present throughout the entire state of NJ, but all of the samples collected in northern NJ were *R. sp. nov.* Our findings therefore have important implications for conservation and geographic range delimitation for not only *R. sp. nov.*, but also *R. sphenoccephala*, which until now has likely been erroneously considered to be part of the fauna of NY and northern NJ.

We strongly suspect that *R. sp. nov.* also occurred on Long Island based on historic descriptive literature and photographs (Overton, 1914a, 1914b; Villani, 1997). Leopard frogs were once abundant on Long Island (Latham, 1971) but are now presumed extirpated (Kiviat, 2010; Feinberg et al., unpublished data). The samples that we analyzed from our field collections on Long Island came from recently metamorphosed tadpoles that our genetic data indicated are *R. palustris*. *Rana palustris* is still common in many central and eastern Long Island localities, and tadpoles and recent metamorphs of this species can be morphologically very similar to leopard frogs. The most recent verified photograph of a live leopard frog on Long Island was taken between 1994 and 1995 (Villani, 1997; Villani, pers. comm.). The historical and current status of leopard frogs on Long Island reflects a distressing trend throughout this region of rapid decline of leopard frog populations (Lannoo, 2005).

The geographic range of *R. sp. nov.* is very small and likely contains only a relatively small number of individual frogs. Amphibians are sensitive to small changes in their environment, and geographically restricted species with few individuals have a reduced chance for survival in the face of rapid climate change, pesticides, and disease (Lande, 1988). *Rana sp. nov.* potentially faces all of these threats, as the pesticide atrazine (Hayes et al.,

2002, 2003, 2010), the fungus *Batrachochytrium dendrobatidis* (Morell, 1999; Bradley et al., 2002; Stuart et al., 2004; Greer et al., 2005; Searle et al., 2011), and *Ranavirus* outbreaks (Granoff et al., 1965) have been shown to have adverse effects on leopard frog populations in this and other regions (but see Voordouw et al., 2010).

Future studies should focus on the ecology and population genetics of *R. sp. nov.*, including breeding phenology and call structure, and incorporate more fine-scaled sampling to gain a better understanding of the distribution of, and gene flow among, existing populations. Ongoing additional work (Feinberg et al., unpublished) will address these issues and describe *R. sp. nov.* as a novel species, furthering our understanding of the *R. pipiens* species complex in this region. In light of this new systematic knowledge, the “precautionary principle” (Raffensperger et al., 1999; Georges et al., 2011) suggests that appropriate conservation measures should be considered for immediate implementation at the state and possibly federal levels. The northeastern US is generally viewed as a glacially-impacted region of low diversity compared to the southeastern US (Rissler and Smith, 2010) or California (Rissler et al., 2006), and thus this region has received relatively less scrutiny and study in recent decades compared to regions that are believed to harbor higher overall diversity (but see *Pseudacris kalmi* Lemmon et al., 2007). However, urban environments such as the northeastern US have been shown to be detrimental to anuran populations, primarily due to habitat fragmentation and isolation, road mortality, and contamination (Findlay and Houlihan, 1997; Hitchings and Beebe, 1997; Knutson et al., 1999; Gibbs et al., 2005). It is therefore likely that species endemic to the Northeast require swift management attention to preserve what biodiversity still remains in the region. Our study revealed a new leopard frog species in the midst of this highly developed region of the US, suggesting that the densely populated Northeast still harbors cryptic biodiversity that remains to be discovered.

#### Acknowledgments

We are indebted to R.C. Thomson, P.Q. Spinks, and G.B. Pauly for bioinformatics help and comments. We would also like to thank M.D. Schlesinger, B. Zarate, and J. Victoria for collecting tissue samples and providing comments, G.J. Watkins-Colwell at the Yale Peabody Museum for providing tissue samples, and J.M. Burnley, F.C. Schlauch, and E. Klaastad for providing comments and valuable natural history information. We thank J.J. Apodaca and G.B. Pauly

for designing primers. We also thank two anonymous reviewers for suggestions that improved this manuscript. Genetics laboratory work was conducted at U.C. Davis, and all research was conducted under appropriate state collecting permits for New York (Collect or Possess, #969, to M.D. Schlessinger) and New Jersey (Scientific Collecting Permit to B. Zarate). Funding for genetic analyses was provided by the National Science Foundation DEB-0817042 to H.B. Shaffer, and the U.C. Davis Agricultural Experiment Station. Additional funding was provided by the New York State Biodiversity Research Institute, Long Island Community Foundation, Rutgers University, Brookhaven National Laboratory, and the Foundation for Ecological Research in the Northeast to J.A. Feinberg.

## Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ympmv.2012.01.021.

## References

- Bradley, G.A., Rosen, P.C., Sredl, M.J., Jones, T.R., Longcore, J.E., 2002. Chytridiomycosis in native Arizona frogs. *J. Wildl. Dis.* 38, 206–212.
- Brown, L.E., 1973. Speciation in the *Rana pipiens* complex. *Am. Zool.* 13, 73–79.
- Brown, L.E., Smith, H.M., Funk, R.S., 1977. Request for the conservation of *Rana sphenocephala* Cope, 1886, and the suppression of *Rana utricularius* Harlan, 1826 and *Rana virescens* Cope, 1889 (Amphibia: Salientia). *Bull. Zool. Nomencl.* 33, 195–203.
- Brown, L.E., Smith, H.M., Funk, R.S., 1990. *Rana sphenocephala* Cope, 1886 (Amphibia, Anura): proposed precedence over *Rana utricularius* Harlan, 1826. *Bull. Zool. Nomencl.* 47, 283–285.
- Carey, C., Cohen, N., Rollins-Smith, L., 1999. Amphibian declines: an immunological perspective. *Dev. Comp. Immunol.* 23, 459–472.
- Conant, R., Collins, J.T., 1998. *A Field Guide to Reptiles & Amphibians: Eastern and Central North America*. Houghton Mifflin Harcourt, New York.
- Davis, A.K., Yabsley, M.J., Keel, M.K., Maerz, J.C., 2007. Discovery of a novel alveolate pathogen affecting southern leopard frogs in Georgia: description of the disease and host effects. *EcoHealth* 4, 310–317.
- de Queiroz, K., 2007. Species concepts and species delimitation. *Syst. Biol.* 56, 879–886.
- Di Candia, M.R., Routman, E.J., 2007. Cytonuclear discordance across a leopard frog contact zone. *Mol. Phylogenet. Evol.* 45, 564–575.
- Ditmars, R.L., n.d. *The Mammals, Reptiles and Amphibians of Westchester County: A Complete Check-list*. Unpublished Manuscript, Wildlife Conservation Society Archives, Bronx, NY.
- Drummond, A.J., Ashton, B., Buxton, S., Cheung, M., Cooper, A., Duran, C., Field, M., Heled, J., Kearse, M., Markowitz, S., Moir, R., Stones-Havas, S., Sturrock, S., Thierer, T., Wilson, A., 2011. Geneious v5.4. <<http://www.geneious.com>>.
- Evanno, G., Regnaut, S., Goudet, J., 2005. Detecting the number of clusters of individuals using the software structure: a simulation study. *Mol. Ecol.* 14, 2611–2620.
- Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Res.* 10, 564–567.
- Falush, D., Stephens, M., Pritchard, J.K., 2003. Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* 164, 1567–1587.
- Findlay, C.S., Houlahan, J., 1997. Anthropogenic correlates of species richness in southeastern Ontario wetlands. *Conserv. Biol.* 11, 1000–1009.
- Frost, D.R., Grant, T., Faivovich, J., Bain, R.H., Haas, A., Haddad, C.F.B., De Sa, R.O., Channing, A., Wilkinson, M., Donnellan, S.C., Raxworthy, C.J., Campbell, J.A., Blotto, B.L., Moler, P., Drewes, R.C., Nussbaum, R.A., Lynch, J.D., Green, D.M., Wheeler, W.C., 2006. The amphibian tree of life. *Bull. Am. Mus. Nat. Hist.* 297, 1–370.
- Frost, D.R., Grant, T., Faivovich, J., Bain, R.H., Haas, A., Haddad, C.F.B., De Sa, R.O., Channing, A., Wilkinson, M., Donnellan, S.C., Raxworthy, C.J., Campbell, J.A., Blotto, B.L., Moler, P., Drewes, R.C., Nussbaum, R.A., Lynch, J.D., Green, D.M., Wheeler, W.C., 2008. Is *The Amphibian Tree of Life* really fatally flawed? *Cladistics* 24, 385–395.
- Frost, D.R., McDiarmid, R.W., Mendelson III, J.R., 2009. Response to the *Point of View* of Gregory B. Pauly, David M. Hillis, and David C. Cannatella, by the anuran subcommittee of the SSAR/HL/ASIH scientific and standard English names list. *Herpetologica* 65, 136–153.
- Funk, W.C., Pearl, C.A., Draheim, H.M., Adams, M.J., Mullins, T.D., Haig, S.M., 2008. Range-wide phylogeographic analysis of the spotted frog complex (*Rana luteiventris* and *Rana pretiosa*) in northwestern North America. *Mol. Phylogenet. Evol.* 49, 198–210.
- Georges, A., Spencer, R.-J., Welsh, M., Shaffer, H.B., Walsh, R., Zhang, X., 2011. Application of the precautionary principle to taxa of uncertain status: the case of the Bellinger River turtle. *Endanger. Species Res.* 14, 127–134.
- Gibbs, J.P., Whiteleather, K.K., Schueler, F.W., 2005. Changes in frog and toad populations over 30 years in New York State. *Ecol. Appl.* 15, 1148–1157.
- Granoff, A., Came, P.E., Rafferty Jr., K.A., 1965. The isolation and properties of viruses from *Rana pipiens*: their possible relationship to the renal adenocarcinoma of the leopard frog. *Ann. N.Y. Acad. Sci.* 126, 237–255.
- Greer, A.L., Berrill, M., Wilson, P.J., 2005. Five amphibian mortality events associated with ranavirus infection in south central Ontario, Canada. *Dis. Aquat. Org.* 67, 9–14.
- Guindon, S., Gascuel, O., 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* 52, 696–704.
- Hayes, T.B., Collins, A., Lee, M., Mendoza, M., Noriega, N., Stuart, A.A., Vonk, A., 2002. Hermaphroditic, demasculinized frogs after exposure to the herbicide atrazine at low ecologically relevant doses. *Proc. Natl. Acad. Sci. U.S.A.* 99, 5476–5480.
- Hayes, T., Haston, K., Tsui, M., Hoang, A., Haeffele, C., Vonk, A., 2003. Atrazine-induced hermaphroditism at 0.1 ppb in American leopard frogs (*Rana pipiens*): laboratory and field evidence. *Environ. Health Perspect.* 111, 568–575.
- Hayes, T.B., Khoury, V., Narayan, A., Nazir, M., Park, A., Brown, T., Adame, L., Chan, E., Buchholz, D., Stueve, T., Gallipeau, S., 2010. Atrazine induces complete feminization and chemical castration in male African clawed frogs (*Xenopus laevis*). *Proc. Natl. Acad. Sci. U.S.A.* 107, 4612–4617.
- Hekkala, E.R., Saumure, R.A., Jaeger, J.R., Herrmann, H.-W., Sredl, M.J., Bradford, D.F., Drabek, D., Blum, M.J., 2011. Resurrecting an extinct species: archival DNA, taxonomy, and conservation of the Vegas Valley leopard frog. *Conserv. Genet.* doi:10.1007/s10592-011-0229-6.
- Hillis, D.M., 1988. Systematics of the *Rana pipiens* complex: puzzle and paradigm. *Annu. Rev. Ecol. Syst.* 19, 39–63.
- Hillis, D.M., Wilcox, T.P., 2005. Phylogeny of the New World true frogs (*Rana*). *Mol. Phylogenet. Evol.* 34, 299–314.
- Hillis, D.M., Frost, J.S., Wright, D.A., 1983. Phylogeny and biogeography of the *Rana pipiens* complex: a biochemical evaluation. *Syst. Zool.* 32, 132–143.
- Hitchings, S.P., Beebe, T.J., 1997. Genetic restructuring as a result of barriers to gene flow in urban *Rana temporaria* (common frog) populations: implications for biodiversity conservation. *Heredity* 79, 117–127.
- Hoffman, E.A., Blouin, M.S., 2004. Evolutionary history of the northern leopard frog: reconstruction of phylogeny, phylogeography, and historical changes in population demography from mitochondrial DNA. *Evolution* 58, 145–159.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogeny. *Bioinformatics* 17, 754–755.
- Jaeger, J.R., Riddle, B.R., Jennings, R.D., Bradford, D.F., McEachran, J.D., 2001. Rediscovering *Rana onca*: evidence for phylogenetically distinct leopard frogs from the border region of Nevada, Utah, and Arizona. *Copeia* 2001, 339–354.
- Kauffeld, C.F., 1937. The status of the leopard frogs, *Rana brachycephala* and *Rana pipiens*. *Herpetologica* 1, 84–87.
- Kiviat, E., 2010. Frog Call Surveys in An Urban Wetland Complex, the Hackensack Meadowlands New Jersey, in 2006. *Urban Habitats* 6. <[http://urbanhabitats.org/v06n01/frogcallsurveys\\_full.html](http://urbanhabitats.org/v06n01/frogcallsurveys_full.html)>.
- Klemens, M.W., 1993. The amphibians and reptiles of Connecticut and adjacent regions. *State Geol. Nat. Hist. Surv. Conn.* 112, 134–140.
- Klemens, M.W., Kiviat, E., Schmidt, R.E., 1987. Distribution of the northern leopard frog, *Rana pipiens*, in the lower Hudson and Housatonic river valleys. *Northeast. Environ. Sci.* 6, 99–101.
- Knutson, M.G., Sauer, J.R., Olsen, D.A., Mossman, M.J., Hemesath, L.M., Lannoo, M.J., 1999. Effects of landscape composition and wetland fragmentation on frog and toad abundance and species richness in Iowa and Wisconsin, U.S.A. *Conserv. Biol.* 13, 1437–1446.
- Köhler, J., Vieites, D.R., Bonett, R.M., García, F.H., Glaw, F., Steinke, D., Vences, M., 2005. New amphibians and global conservation: a boost in species discoveries in a highly endangered vertebrate group. *BioScience* 55, 693–696.
- Kubatko, L.S., Degnan, J.H., 2007. Inconsistency of phylogenetic estimates from concatenated data under coalescence. *Syst. Biol.* 56, 17–24.
- Lande, R., 1988. Genetics and demography in biological conservation. *Science* 241, 1455–1460.
- Lannoo, M. (Ed.), 2005. *Amphibian Declines: The Conservation Status of United States Species*. University of California Press, USA.
- Lannoo, M., 2008. *Malformed Frogs: The Collapse of Aquatic Ecosystems*. University of California Press, USA.
- Latham, R., 1971. The leopard frog on eastern Long Island. *Engelhardtia* 4, 58.
- Lemmon, E.M., Lemmon, A.R., Collins, J.T., Lee-Yaw, J.A., Cannatella, D.C., 2007. Phylogeny-based delimitation of species boundaries and contact zones in the trilling chorus frogs (*Pseudacris*). *Mol. Phylogenet. Evol.* 44, 1068–1082.
- Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25, 1451–1452.
- Moore, J.A., 1975. *Rana-pipiens*: the changing paradigm. *Am. Zool.* 15, 837–849.
- Morell, V., 1999. Are pathogens felling frogs? *Science* 284, 728–731.
- Newman, C.E., Rissler, L.J., 2011. Phylogeographic analyses of the southern leopard frog: the impact of geography and climate on the distribution of genetic lineages vs. subspecies. *Mol. Ecol.* 20, 5295–5312.
- Niemiller, M.L., Near, T.J., Fitzpatrick, B.M., 2011. Delimiting species using multilocus data: diagnosing cryptic diversity in the southern cavefish, *Typhlichthys subterraneus* (Teleostei: Amblyopsidae). *Evolution*. doi:10.1111/j.1558-5646.2011.01480.x.
- Oláh-Hemmings, V., Jaeger, J.R., Sredl, M.J., Schlaepfer, M.A., Jennings, R.D., Drost, C.A., Bradford, D.F., Riddle, B.R., 2010. Phylogeography of declining relict and lowland leopard frogs in the desert Southwest of North America. *J. Zool.* 280, 343–354.

- O'Meara, B.C., 2010. New heuristic methods for joint species delimitation and species tree inference. *Syst. Biol.* 59, 59–73.
- Overton, F., 1914a. Long Island fauna and flora III: the frogs and toads. *Mus. Brooklyn Inst. Arts Sci. Bull.* 2, 21–53.
- Overton, F., 1914b. The frogs and toads of Long Island. *Brooklyn Mus. Q.* 1, 30–38.
- Pace, A.E., 1974. Systematic and biological studies of the leopard frogs (*Rana pipiens* complex) of the United States. *Univ. Mich. Mus. Zool.* 148, 1–140.
- Pauly, G.B., Hillis, D.M., Cannatella, D.C., 2009. Taxonomic freedom and the role of official lists of species names. *Herpetologica* 65, 115–128.
- Posada, D., 2008. JModelTest: phylogenetic model averaging. *Mol. Biol. Evol.* 25, 1253–1256.
- Pritchard, J.K., Stephens, M., Donnelly, P., 2000. Inference of population structure using multilocus genotype data. *Genetics* 155, 945–959.
- Pytel, B.A., 1986. Biochemical systematics of the eastern North American frogs of the genus *Rana*. *Herpetologica* 42, 273–282.
- Raffensperger, C., Tickner, J., Jackson, W., 1999. Protecting Public Health and the Environment: Implementing the Precautionary Principle. Island Press, Washington, DC.
- Rambaut, A., Drummond, A.J., 2007. *Tracer v.1.4*. <<http://beast.bio.ed.ac.uk/Tracer>>.
- Rissler, L.J., Smith, W.H., 2010. Mapping amphibian contact zones and phylogeographical break hotspots across the United States. *Mol. Ecol.* 19, 5404–5416.
- Rissler, L.J., Hijmans, R.J., Graham, C.H., Moritz, C., Wake, D.B., 2006. Phylogeographic lineages and species comparisons in conservation analyses: a case study of California herpetofauna. *Am. Nat.* 167, 655–666.
- Ronquist, F., Huelsenbeck, J.P., 2003. MRBAYES3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19, 1572–1574.
- Schlauch, F.C., 1971. The subspecific status of leopard frogs of a region in the Pine Barrens of Long Island. *Engelhardtia* 4, 47–49.
- Schlauch, F.C., 1978. Literature review: endangered amphibians and reptiles. *Pitch Pine Nat.* 4, 5–6.
- Searle, C.L., Gervasi, S.S., Hua, J., Hammond, J.I., Relyea, R.A., Olson, D.H., Blaustein, A.R., 2011. Differential host susceptibility to *Batrachochytrium dendrobatidis*, an emerging amphibian pathogen. *Conserv. Biol.* doi:10.1111/j.1523-1739.2011.01708.x.
- Shaffer, H.B., Thomson, R.C., 2007. Delimiting species in recent radiations. *Syst. Biol.* 56, 896–906.
- Shaffer, H.B., Fellers, G.M., Voss, S.R., Oliver, J.C., Pauly, G.B., 2004. Species boundaries, phylogeography and conservation genetics of the red-legged frog (*Rana aurora/draytonii*) complex. *Mol. Ecol.* 13, 2667–2677.
- Sites Jr., J.W., Marshall, J.C., 2004. Operational criteria for delimiting species. *Annu. Rev. Ecol. Syst.* 35, 199–227.
- Stamatakis, A., 2006. RAXML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22, 2688–2690.
- Stamatakis, A., Hoover, P., Rougemont, J., 2008. A rapid bootstrap algorithm for the RAXML web servers. *Syst. Biol.* 57, 758–771.
- Stephens, M., Donnelly, P., 2003. A comparison of Bayesian methods for haplotype reconstruction. *Am. J. Hum. Genet.* 73, 1162–1169.
- Stephens, M., Smith, N.J., Donnelly, P., 2001. A new statistical method for haplotype reconstruction from population data. *Am. J. Hum. Genet.* 68, 978–989.
- Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L., Waller, R.W., 2004. Status and trends of amphibian declines and extinctions worldwide. *Science* 306, 1783–1786.
- Vähä, J.P., Primmer, C.R., 2005. Efficiency of model-based Bayesian methods for detecting hybrid individuals under different hybridization scenarios and with different numbers of loci. *Mol. Ecol.* 15, 63–72.
- Villani, R., 1997a. Long Island: A Natural History. Harry N. Abrams, Inc., NY.
- Voordouw, M.J., Adama, D., Houston, B., Govindarajulu, P., Robinson, J., 2010. Prevalence of the pathogenic chytrid fungus, *Batrachochytrium dendrobatidis*, in an endangered population of northern leopard frogs, *Rana pipiens*. *BMC Ecol.* 10. doi:10.1186/1472-6785-10-6.
- Yeaton, S.C., Jr., 1968. The Amphibia of Long Island. Sanctuary Summer, pp. 2–19.
- Zug, G.R., Wassersug, R., Uzzell, T., Brown, L.E., Smith, H.M., Funk, R.S., 1982. Comments on the proposed conservation of *Rana sphenoccephala* Cope, 1886. *Bull. Zool. Nomencl.* 39, 80–90.